



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

# ON A NEW CURVE FOR THE TRISECTION OF AN ANGLE.

BY DR. WILLIAM HILLHOUSE.

THE following method for the trisection of an angle from  $0$  to  $360^\circ$ , is believed to be new. The eq'n to the curve has not been found in any book relating to curves that has come to the notice of the writer, nor has its mechanical description ever been seen. It is now offered to those interested in such matters, as a new contribution to the subject, and if it should excite any interest in them, or in any way prove of value in geometry, the object of the writer will be attained.

$ABCD$ , Fig. 1, represents a jointed parallelogram ( $AC$ ,  $FE$  and  $BD$  are supposed to be infinite). Into  $CD$  is fixed firmly the piece  $FE$ , whose edge  $FE$  is at right angles to  $CD$ . This piece  $FE$  is about one-third the thickness of  $CD$  to admit of its passing freely under the bar  $AC$ , which is elevated on the under side for that purpose. The point  $F$  is also equidistant from the inner edges of  $AC$  and  $BD$ . If the bar  $BD$  be fixed on a plane, and the bar  $AC$  be depressed, as indicated by the dotted lines in the figure,  $AC$  will always remain parallel to  $BD$ , and the bar  $FE$  will incline to the left, as  $fe$ , and intersect the edge of  $AC$ . The p't of intersection of the inner edges of  $AC$  and  $FE$  will trace out (by applying a pencil at the point) a locus, represented in part by the line  $bpc$ , which will be of the fourth order, having four infinite branches,  $AH$ ,  $AG$ ,  $BN$ ,  $BK$  (Fig. 2). The inner edge of  $AC$ , when it is parallel to  $EF$  will be an asymptote to the curve. The eq'n of the curve may be found as follows, by refer'g to Fig. 2.




Fig 1.



$$\therefore z^2 = \frac{4a^2x^2}{(r-x)^2} = x^2 + \frac{x^2y^2}{(r-x)^2},$$


$$\text{or } 4a^2 = (r-x)^2 + y^2. \quad (1)$$

Substituting for  $r$  in (1), and reducing, we have finally

$$x^2 = \frac{(2a^2 - y^2)^2}{4a^2 - y^2}$$

for the equation of the locus.

To trisect any given angle, draw a straight line (Fig. 3)  $BOX$  and a straight line  $bBD$  at right angles to it. From  $B$  lay off  $BO = a$ ; with  $O$  as centre and distance  $BO$ , describe a circle  $fBG$ , and also from the same point a circle with radius  $OC = 2a$ . Let  $BOX$  be the initial line from which the given angle is measured, passing around to the left; let  $HOX$  be the angle to be trisected. From  $O$  draw  $OKN$  at right angles to  $HO$ , and place the parallelogram so that the inner edge of  $CD$  (Fig. 1) is on the line  $HO$ , and the inner edge of  $BD$  (Fig. 1) on  $OKN$ ; depress the side  $AC$  of the parallelogram so that the intersection of the lines  $AC$  and  $FE$  shall be on the line  $DBb$ , say the point  $b$ ; join  $bO$ , and from the point  $b$  draw  $bf$  tangent to the circle  $BGf$ ; from  $O$  through  $f$  draw  $OfC$ , which will be perpendicular to  $bf$ , and join  $bC$ . Now the angle  $OKB = CbB = HOX$ ,



and in the right angled triangles  $bBO$ ,  $bfO$ , the side  $BO =$  the side  $Of$ , since they are radii of the same circle, and the side  $bO$  common, therefore the other angles are equal, that is  $O\hat{b}B = O\hat{b}f$ ; also in the right triangles  $bfO$ ,  $bfC$ , the side  $Of =$  the side  $fC$  by construction and the side  $bf$  is common, therefore the angle  $O\hat{b}f =$  the angle  $C\hat{b}f$ . But  $O\hat{b}f$  was shown to be  $= O\hat{b}B$ ; therefore  $C\hat{b}f = O\hat{b}B$ , that is, the three angles are equal to each other. But the three angles  $C\hat{b}f$ ,  $f\hat{b}O$ ,  $O\hat{b}B$ , make up the angle  $C\hat{b}B = HOX$  the given angle.

Now in the right angled triangles  $bfC$  and  $OIC$  the angle  $bCO$  is common to the two triangles, therefore the remaining angles are equal, that is, the angle  $COI =$  the angle  $C\hat{b}f$ ; but  $C\hat{b}f$  has been shown to be equal to  $O\hat{b}B$ , therefore  $COI = O\hat{b}B$ ; but  $O\hat{b}B$  is one-third of  $C\hat{b}B$  or  $HOX$ , therefore  $HOC =$  one-third  $HOX$ .

WM. HILLHOUSE.

New Haven, Conn., Jan., 1878.

SOLUTION OF PROB. 405 BY PROF. C. A. VAN VELZER.—To fix the idea take the determinant of the fourth order

$$\begin{vmatrix} a_1 & b_1 & c_1 & d_1 \\ a_2 & b_2 & c_2 & d_2 \\ a_3 & b_3 & c_3 & d_3 \\ a_4 & b_4 & c_4 & d_4 \end{vmatrix} = 0.$$

To the second column add  $l_1$  times the first,  $m_1$  times the third, and  $n_1$  times the fourth, where  $l_1, m_1, n_1$  are chosen to satisfy the equations

$$\begin{aligned} (l_1-1)a_1 + m_1c_1 + n_1d_1 + b_1 &= 0, \\ (l_1-1)a_2 + m_1c_2 + n_1d_2 + b_2 &= 0, \\ (l_1-1)a_3 + m_1c_3 + n_1d_3 + b_3 &= 0. \end{aligned}$$

These three equations are sufficient to determine the values of  $(l_1-1), m_1, n_1$ , but if to these we add a fourth

$$(l_1-1)a_4 + m_1c_4 + n_1d_4 + b_4 = 0$$

these *four* equations form a consistent set in  $(l_1-1), m_1, n_1$ , since the determinant of the coefficients (viz. the original determinant) vanishes.

We see that by this first transformation the determinant reduces to

$$\begin{vmatrix} a_1 & a_1 & c_1 & d_1 \\ a_2 & a_2 & c_2 & d_2 \\ a_3 & a_3 & c_3 & d_3 \\ a_4 & a_4 & c_4 & d_4 \end{vmatrix}$$

Now to the second row of this determinant add  $l_2$  times the first,  $m_2$  times the third and  $n_2$  times the fourth, where  $l_2, m_2, n_2$  satisfy the equ'n's